

<b>Time Point</b>	<b>Time/Duration</b>	<b>Task/Description</b>
T0	--	Orthodontist makes dx of jaw problem likely needing surgery Refer patient to Dr. Steinbacher
T1a	early	Visit w/ Dr. Steinbacher – Jaw assessment
T1b	6-18 mo	PreSurgical Orthodontics
T2a	6-10 mo in to orthodontic tx	Choose a surgery date (patient/family based on orthodontist input as to when ready; and Dr. Steinbacher schedule) Orthodontist gauge/tell patient and Dr. Steinbacher when presurgical orthodontics will be completed
T2b	4-6 weeks before surgery	Orthodontist places surgical wire, rectangular, passive Hooks placed at that time, or the week before surgery
T2c	8-14 days before surgery	Dr. Steinbacher preop visit; obtain CBCT, dental casts
T2d	7-12 days before surgery	Final occlusion set – with input from orthodontist 3D occlusion fused with CBCT to prepare for planning
T2e	6-7 days before surgery	3D planning session – orthodontist can join Splints 3D printed and sent for surgery
T3	--	Surgery
T4	7-9 days after surgery	Postop visit with Dr. Steinbacher Elastic loosened
T5	3-6 weeks after surgery	Orthodontist first postop visit At 6 weeks can dictate how elastics used, or if they are needed
T5a	3-6 weeks after surgery	Hawley type palatal splint removed – if a segmental Le Fort Either Dr. Steinbacher or orthodontist can remove this Please place a continuous rigid arch-wire to span segments within 1 day of removal Or place a clear aligner / retainer based on new maxillary width within 1 day of removal (if Invisalign patients)
T5b	6-12 weeks after surgery	Remove TADs if they are no longer needed Dr. Steinbacher or orthodontist can remove these
T6	4-6 weeks + after surgery For 4-12 mo	PostSurgical orthodontic phase
T6a	2-4 mo + after surgery	Place dental implants if they are needed / planned
T6a	2-4 mo + after surgery	Perform Rhinoplasty / nasal procedure if indicated / needed / planned
T7	Several months after surgery	Braces off – retention phase Completion of treatment

\*In cases of sleep apnea we will obtain a postoperative sleep study usually 4-6 mo after jaw surgery

\*In cleft palate cases, speech surgery may be needed / planned / considered after orthognathic surgery

\*in some cases hardware is removed after orthognathic surgery – the soonest we usually would do this is 3-4 months after surgery

BLUE ink = ORTHODONTIST VISIT / ROLE

Green ink = Dr. Steinbacher